



PTO/SB/ 42 (6-95)

Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Sheet 1 of 9

37 CFR 1.501b

INFORMATION DISCLOSURE
CITATION IN A PATENT

(Use several sheets if necessary)

Docket Number: 04340-PCT-PA (0148.0041)

Serial Number: 10/516,406

Applicant: PETRONIS

Issue Date:

Group Art Unit: 1647/1634

U.S. PATENT DOCUMENTS

Examiner Initial	Document Number	Date	Name	Class	Subclass	Filing Date if Appropriate
/CM/	4,683,195	07/28/1987	Mullis et al	435	6	
↓	4,683,202	07/28/1987	Mullis	435	91	
↓	4,800,159	01/24/1989	Mullis et al	435	172.3	
↓	5,871,917	02/16/1999	Duffy	435	6	

	Document Number	Date	Country	Class	Subclass	Translation	
						Yes	No
/CM/	US98/13818	07/02/1998	PCT				

OTHER DOCUMENTS (Including Author, Title, Date, Pertinent Pages, Etc.)

	87 references. Please see attachment.

EXAMINER

/Carla Myers/

Date Considered

06/11/2007

Burden Hour Statement: This form is estimated to take 2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

Other Documents

Alitalo T, Tiihonen J, Hakola P, de la Chapelle A (1988): Molecular characterization of a Y;15 translocation segregating in a family. Hum Genet 79:29-35.

Altschul et al., J. Mol. Biol. (1990) 215:403-10

Bassett AS, Chow EW, Waterworth DM, Brzustowicz L (2001): Genetic insights into schizophrenia. Can J. Psychiatry 46:131-7.

Bennett RL, Karayiorgou M, Sobin CA, Norwood TH, Kay MA (1997): Identification of an interstitial deletion in an adult female with schizophrenia, mental retardation, and dysmorphic features: further support for a putative schizophrenia-susceptibility locus at 5q21-23.1. Am J Hum Genet 61:1450-4.

Berrettini W (2002): Review of bipolar molecular linkage and association studies. Curr Psychiatry Rep 4:124-9.

Berrettini WH 2000a): Are schizophrenic and bipolar disorders related? A review of family and molecular studies. Biol Psychiatry 48:531-8.

Berrettini WH 2000b): Susceptibility loci for bipolar disorder; overlap with inherited vulnerability to schizophrenia. Biol Psychiatry 47:245-51.

Bray NJ, Owen MJ. Searching for schizophrenia genes. Trends Mol Med. 2001; 7(4):169-74.

Brzustowicz LM, Hodgkinson KA, Chow EW, Honer WG, Bassett AS. Location of a major susceptibility locus for familial schizophrenia on chromosome 1q21-q22. Science 2000 Apr 28;288(5466):678-82.

Camp NJ, Neuhausen SL, Tiobech J, Polloi A, Coon H, Myles-Worsley M (2001): Genomewide multipoint linkage analysis of seven extended Palauan pedigrees with schizophrenia, by a Markov-chain Monte Carlo method. Am J Hum Genet 69:1278-89.

Cardno AG, Rijsdijk FV, Sham PC, Murray RM, McGuffin P (2002): A twin study of genetic relationships between psychotic symptoms. Am J Psychiatry 159:539-45.

Chow EW, Bassett AS, Weksberg R. Velo-cardio-facial syndrome and psychotic disorders: implications for psychiatric genetics. *Am J Med Genet* 1994;54(2):107-12

DeLisi LE, Shaw SH, Crow TJ, et al (2002): A genome-wide scan for linkage to chromosomal regions in 382 sibling pairs with schizophrenia or schizoaffective disorder. *Am J Psychiatry* 159:803-12.

Detera-Wadleigh SD, Badner JA, Berrettini WH, et al (1999): A high-density genome scan detects evidence for a bipolar-disorder susceptibility locus on 13q32 and other potential loci on 1q32 and 18p11.2. *Proc Natl Acad Sci U S A* 96:5604-9

Detera-Wadleigh SC, Badner JA, Goldin LR, et al (1996): Affected-sib-pair analyses reveal support of prior evidence for a susceptibility locus for bipolar disorder, on 21q. *Am J Hum Genet* 58:1279-85.

Egeland et al. (1987) *Nature* 325: 783-787.

Evans KL, Brown J, Shibasaki Y, et al (1995): A contiguous clone map over 3 Mb on the long arm of chromosome 11 across a balanced translocation associated with schizophrenia. *Genomics* 28:420-8.

Ewald H, Wang AG, Vang M, Mors O, Nyegaard M, Kruse TA (1999): A haplotype-based study of lithium responding patients with bipolar affective disorder on the Faroe Islands. *Psychiatr Genet* 9:23-34.

Fan et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6223-27.

Faraone SV, Matise T, Svrakic D, et al (1998): Genome scan of European-American schizophrenia pedigrees: results of the NIMH Genetics Initiative and Millennium Consortium. *Am J Med Genet* 81:290-5.

Feinberg, A. P. and Vogelstein, B. (1983) Hypomethylation of ras oncogenes in primary human cancers. *Biochem Biophys Res Commun* 111, 47-54.

Feinberg, A. P., Gehrke, C. W., Kuo, K. C., and Ehrlich, M. (1988). Reduced genomic 5-methylcytosine content in human colonic neoplasia. *Cancer Res* 48, 1159-61.

Filippova GN, Thienes CP, Penn BH, et al (2001): CTCF-binding sites flank CTG/CAG repeats and form a methylation-sensitive insulator at the DM1 locus. *Nat Genet* 28:335-43.

Florl AR, Franke KH, Niederacher D, Gerharz CD, Seifert HH, Schulz WA. DNA methylation and the mechanisms of CDKN2A inactivation in transitional cell carcinoma of the urinary bladder. *Laboratory Investigation*, 80(10):1513-22, 2000.

Fremmer NB, Reus VI, Escamilla MA, et al (1996): Genetic mapping using haplotype, association and linkage methods suggests a locus for severe bipolar disorder (BPI) at 18q22-q23. *Nat Genet* 12:436-41.

Gama-Sosa, M.A., Slagel, V. A., Trewyn, R. W., Oxenhandler, R., Kuo, K. C., Gehrke, C. W., and Ehrlich, M. (1983). The 5-methylcytosine content of DNA from human tumors. *Nucleic Acids Res* 11, 6883-94.

Gershon ES. Bipolar illness and schizophrenia as oligogenic diseases: implications for the future. *Biol Psychiatry*. 2000 Feb 1; 47(3):240-4.

Gill M, Vallada H, Collier D, et al (1996): A combined analysis of D22S278 marker alleles in affected sib-pairs: support for a susceptibility locus for schizophrenia at chromosome 22q12. Schizophrenia Collaborative Linkage Group (Chromosome 22). *Am J Med Genet* 67:40-5.

Gonzalzo, M.L. and Jones, P.A. (1997) Mutagenic and epigenetic effects of DNA methylation. *Mutat. Res.* 386(2), 107-18.

H. Suzuki et al., 6 *Nature Genet* 332-33 (1994) (please note that the specification incorrectly lists this reference as H. Suzuki et al., 7 *Nature Genet* 432-38 (1994))

Hedenfalk I, Duggan D, Chen Y, et al (2001): Gene-expression profiles in hereditary breast cancer. *N Engl J Med* 344:539-48.

Henikoff S, Matzke MA Exploring and explaining epigenetic effects. *Trends Genet* 1997; 13(8):293-5.

Hernandez I, Sokolov BP. Abnormalities in 5-HT_{2A} receptor mRNA expression in frontal cortex of chronic elderly schizophrenics with varying histories of neuroleptic treatment. *J Neurosci Res.* 2000; 59(2):218-25.

Issa J. P., et al. (1994) *Nature Genetics* 7:536-40.

Johnston-Wilson NL, Sims CD, Hofmann JP, et al (2000): Disease-specific alterations in frontal cortex brain proteins in schizophrenia, bipolar disorder, and major depressive disorder. The Stanley Neuropathology Consortium. *Mol Psychiatry* 5:242-9.

Joo EJ, Lee JH, Cannon TD, Price RA (1999): Possible association between schizophrenia and a CAG repeat polymorphism in the spinocerebellar ataxia type 1 (SCA1) gene on human chromosome 6p23. *Psychiatr Genet* 9:7-11.

Jurgens B, Schmitz-Drager BJ, Schulz WA. Hypomethylation of L1 LINE sequences prevailing in human urothelial carcinoma. *Cancer Research*, 56(24):5698-703, 1996.

Karlsson H, Bachmann S, Schroder J, McArthur J, Torrey EF, Yolken RH (2001): Retroviral RNA identified in the cerebrospinal fluids and brains of individuals with schizophrenia. *Proc Natl Acad Sci U S A* 98:4634-9.

Kelsoe JR, Spence MA, Loetscher E, et al (2001): A genome survey indicates a possible susceptibility locus for bipolar disorder on chromosome 22. *Proc Natl Acad Sci U S A* 98:585-90.

Kendler KS, Myers JM, O'Neill FA, et al (2000): Clinical features of schizophrenia and linkage to chromosomes 5q, 6p, 8p, and 10p in the Irish Study of High-Density Schizophrenia Families. *Am J Psychiatry* 157:402-8.

Kwoh et al., 8 *Am. Biotechnol. Lab.* 14 (1990).

Laird et al., *DNA Methylation and Cancer*, 3 *Human Molecular Genetics* 1487, 1488 (1994).

Lapeyre, J. N. and Becker, F. F. (1979). 5-Methylcytosine content of nuclear DNA during chemical hepatocarcinogenesis and in carcinomas which result. *Biochem Biophys Res Commun* 87, 698-705.

Le Corre S, Harper CG, Lopez P, Ward P, Catts S (2000). Increased levels of expression of an NMDAR1 splice variant in the superior temporal gyrus in schizophrenia. *Neuroreport* 11:983-6.

Liang G, et al. (2002) Identification of DNA methylation differences during tumorigenesis by methylation-sensitive arbitrarily primed polymerase chain reaction. *Methods* 27(2): 150-5.

McNeil TF (1995): Perinatal risk factors and schizophrenia: selective review and methodological concerns. *Epidemiol Rev* 17:107-12.

Miniou P, Bourc'his D, Molina Gomes D, Jeanpierre M, Viegas-Pequignot E (1997): Undermethylation of Alu sequences in ICF syndrome: molecular and in situ analysis. *Cytogenet Cell Genet* 77:308-13.

Mirnics K, Middleton FA, Lewis DA, Levitt P (2001): Analysis of complex brain disorders with gene expression microarrays: schizophrenia as a disease of the synapse. Trends Neurosci 24:479-86.

Meises HW, Yang L, Li T, et al (1995): Potential linkage disequilibrium between schizophrenia and locus D22S278 on the long arm of chromosome 22. Am J Med Genet 60:465-7.

Mors O, Mortensen PB, Ewald H (2001): No evidence of increased risk for schizophrenia or bipolar affective disorder in persons with aneuploidies of the sex chromosomes. Psychol Med 31:425-30.

Mowry BJ, Nancarrow DJ (2001): Molecular genetics of schizophrenia. Clin Exp Pharmacol Physiol 28:66-9.

Mujaheed M, Corbex M, Dichtenberg P, et al (2000): Evidence for linkage by transmission disequilibrium test analysis of a chromosome 22 microsatellite marker D22S278 and bipolar disorder in a Palestinian Arab population. Am J Med Genet 96:836-8.

Myles-Worsley M, Coon H, McDowell J, et al (1999): Linkage of a composite inhibitory phenotype to a chromosome 22q locus in eight Utah families. Am J Med Genet 88:544-50.

Nurnberger JI Jr, Foroud T. Genetics of bipolar affective disorder. Curr Psychiatry Rep 2000 Apr;2(2):147-57.

Ogawa, et al., 362 Nature 749-51 (1993).

P. V. Pedone et al., 3 Hum. Mol. Genet. 1117-21 (1994).

Petit J, Boisseau P, Taine L, Gauthier B, Arveiler B (1999): A YAC contig encompassing the 11q14.3 breakpoint of a translocation associated with schizophrenia, and including the tyrosinase gene. Mamm Genome 10:649-52.

Petronis A, Gottesman, II, Crow TJ, et al (2000): Psychiatric epigenetics: a new focus for the new century. Mol Psychiatry 5:342-6.

Petronis A, Paterson AD, Kennedy JL. Schizophrenia: an epigenetic puzzle? Schizophrenia Bulletin 25:4: 639-655, 1999; Petronis A. The genes for major psychosis: aberrant sequence or regulation? Neuropsychopharmacology, 23(1):1-12; 2000.

Petronis A. Human morbid genetics revisited: relevance of epigenetics. Trends Genet. 2001 Mar;17(3):142-6.

Petronis A. The genes for major psychosis: aberrant sequence or regulation? Neuropsychopharmacology, 23(1):1-12; 2000.

Pfeifer GP, et al. (1989) Science 246:810-13.

Popendikyte V, Laurinavicius A, Paterson AD, Macciardi F, Kennedy JL, Petronis A. DNA methylation at the putative promotor region of the human dopamine D2 receptor gene. Neuroreport 1999; 10:1249-55.

Pulver AE, Karayiorgou M, Wolyniec PS, et al (1994): Sequential Strategy to identify a susceptibility gene for schizophrenia: report of potential linkage on chromosome 22q12-q13.1:Part 1. Am J Med Genet 54:36-43.

S. Rainier et al, 55 Cancer Res. 1836-38 (1995).

Rainier et al. 362 Nature 747-49 (1993).

Rakyan VK, Blewitt ME, Druker K, Preis JJ, Whitelaw E (2002): Metastable epialleles in mammals. Trends Genet 18:348-51.

Rice JP, Goate A, Williams JT, et al (1997): Initial genome scan of the NIMH genetics initiative bipolar pedigrees: chromosomes 1, 6, 8, 10, and 12. Am J Med Genet 74:247-53.

Robertson KD and Wolffe AP (2000) DNA methylation in health and disease. Nature Review Genet 1:11-9.

S. Zhan et al., 94 J. Clin. Invest. 445-48 (1994).

Saiki et al. (1988 Science 239:487-491).

Salz and Jost (1989) Proc Natl Acad Sci USA 86:2602-6.

Schizophrenia Collaborative Linkage Group (1998): A transmission disequilibrium and linkage analysis of D22S278 marker alleles in 574 families: further support for a susceptibility locus for schizophrenia at 22q12. Schizophrenia Res 32:115-21.

Schwab SG, Albus M, Hallmayer J, et al (1995a): Evaluation of a susceptibility gene for schizophrenia on chromosome 6p by multipoint affected sib-pair linkage analysis. *Nat Genet* 11:325-7.

Schwab SG, Hallmayer J, Albus M, et al (1998): Further evidence for a susceptibility locus on chromosome 10p14-p11 in 72 families with schizophrenia by nonparametric linkage analysis. *Am J Med Genet* 81:302-7.

Schwab SG, Lerer B, Albus M, et al (1995b): Potential linkage for schizophrenia on chromosome 22q12-q13: a replication study. *Am J Med Genet* 60:436-43.

Sherrington R, Brynjolfsson J, Petursson H, Potter M, Duddleston K, Barraclough B, Wasmuth J, Dobbs M, Gurling H. Localization of a susceptibility locus for schizophrenia on chromosome 5. *Nature*. 1988 Nov 10:336(6195):164-7.

Sinnett D, Richer C, Deragon JM, Labuda D. Alu RNA transcripts in human embryonal carcinoma cells. Model of post-transcriptional selection of master sequences. *Journal of Molecular Biology*. 226(3):689-706, 1992.

St Clair D, Blackwood D, Muir W, Carothers A, Walker M, Spowart G, Gosden C, Evans HJ. Association within a family of a balanced autosomal translocation with major mental illness. *Lancet*. 1990 Jul 7;336(8706):13-6.

Straub RE, MacLean CJ, Martin RB, et al (1998): A schizophrenia locus may be located in region 10p15-p11. *Am J Med Genet* 81:296-301.

Straub RE, MacLean CJ, O'Neill FA, Walsh D, Kendler KS (1997): Support for a possible schizophrenia vulnerability locus in region 5q22-31 in Irish families. *Mol Psychiatry* 2:148-55.

Susser E, Neugebauer R, Hoek HW, et al (1996): Schizophrenia after prenatal famine. Further evidence. *Arch Gen Psychiatry* 53:25-31.

Thaker GK, Carpenter WT Jr. Advances in schizophrenia. *Nat Med*. 2001 Jun;7(6):667-71.

Tsuang MT, Stone WS, Faraone SV. Genes, environment and schizophrenia. *Br J Psychiatry Suppl*. 2001 Apr; 40:s18-24.

Verheyen GR, Villafuerte SM, Del-Favero J, et al (1999): Genetic refinement and physical mapping of a chromosome 18q candidate region for bipolar disorder. *Eur J Hum Genet* 7:427-34.

Wang S, Detera-Wadleigh SD, Coon H, et al (1996): Evidence of linkage disequilibrium between schizophrenia and the SCA1 CAG repeat on chromosome 6p23. Am J Hum Genet 59:731-6.

Xu G-L, Bestor TH. Nature Genetics 17:376-379, 1997.